

the plasma wave and backscattered EM wave are approximately equal, half the energy leaving the pump will go directly into the unstable plasma wave. All of the energy in this electrostatic wave will eventually be locally absorbed by the plasma near the cutoff of the EM wave since the plasma wave is nonpropagating. When the density gradient is weak, most of the energy in the scattered EM wave is trapped in the resonant region of the plasma wave, and therefore a large fraction of this energy will eventually be locally absorbed by the plasma. In stronger density gradients most of the energy in the scattered EM wave will escape from the plasma.

The authors would like to acknowledge helpful conversations with A. Baños, Jr., B. D. Fried, J. M. Kindel, and G. Schmidt.

\*Work partially supported by the U.S. Office of Naval

Research under Grant No. N00014-69-A-0200-4023; the National Science Foundation under Grant No. GP-22817; and the U.S. Atomic Energy Commission under Contract No. AT(04-3)-34 P.A. 157.

<sup>1</sup>M. V. Rosenbluth, Phys. Rev. Lett. **29**, 565 (1972); C. S. Liu, Bull. Amer. Phys. Soc. **17**, 1065 (1972); D. Pesme, G. Laval, and R. Pellat, Phys. Rev. Lett. **31**, 203 (1973).

<sup>2</sup>D. W. Forslund, J. M. Kindel, and E. L. Lindman, Phys. Rev. Lett. **30**, 739 (1973).

<sup>3</sup>J. M. Kindel, private communication.

<sup>4</sup>The inversion of Eq. (1) has been carried out by J. Drake and Y. C. Lee [UCLA Plasma Physics Report No. PPG 156 (unpublished)] by obtaining an exact integral representation for  $E_\omega$  and then expanding the integral.

<sup>5</sup>Equation (4) has been generalized to the case of two-dimensional scattering ( $E_{\omega_0} \cdot \nabla = 0$ ) by Drake and Lee, Ref. 4.

<sup>6</sup>A. Baños, Jr., UCLA Plasma Physics Report No. PPG 124 (to be published).

<sup>7</sup>Drake and Lee, Ref. 4.

## Study of the Bend Elastic Constant near a Smectic-A–Nematic Phase Transition

P. E. Cladis

*Bell Laboratories, Murray Hill, New Jersey 07974*

(Received 16 April 1973; revised manuscript received 27 September 1973)

The temperature dependence of the bend elastic constant of N-*p*-cyanobenzilidene-*p*-*n*-octyloxyaniline near the smectic-A–nematic transition temperature  $T_c$  is found to obey the power law  $(T - T_c^*)^{-\gamma_3}$  with classical values for  $\gamma_3$  in purer samples. Deviations from this value are discussed. Scanning calorimetry indicates that for purer samples the transition is either weakly first order or a  $\lambda$  transition.

Recently, de Gennes<sup>1</sup> has formulated an analogy between the superconducting-normal metal transition and the second-order nematic–smectic-A transition. In the former case, the fluctuations of temporal Cooper pairs in the normal state lead to an increase in the normal-state diamagnetic susceptibility near  $T_c$ .<sup>2</sup> In the latter, the density fluctuations in the nematic phase (which may be described by an order parameter,  $\psi$ ), due to the continual production of evanescent, submicroscopic smectic regions, lead to an increase in the nematic elastic constants<sup>3</sup> of bend ( $K_3$ ) and twist ( $K_2$ ) so that

$$\delta K_i = K_i(T) - (K_i)_0 = (\text{const})_i (T - T_c)^{-\gamma_i}, \quad (1)$$

where  $T_c$  is the nematic–smectic transition temperature,  $(K_i)_0$  is the value without the enhancement due to the order-parameter fluctuations,  $i = 2, 3$ . Here measurements are presented on the temperature dependence of  $K_3(T)$  for CBOOA (N-

*p*-cyanobenzilidene-*p*-*n*-octyloxyaniline). Fitting the data with the functional form of Eq. (1), one determines the four parameters  $(K_i)_0$ ,  $(\text{const})_i$ ,  $\gamma_i$ , and  $T_c$  which best fit the data. This is done by the well-known technique of minimizing the weighted sum of the squares of the difference between the measured and computed values (i.e.,  $\chi^2$ ) for  $K_i(T)$  with respect to the four parameters. In the case of a second-order transition,  $T_c$  may be measured independently and so does not enter as a parameter in the fit. In the case of a first-order transition the  $T_c$  in Eq. (1) should be replaced by  $T_c^* < T_{c, \text{meas}}$  which is to be found.

CBOOA was chosen because McMillan<sup>4</sup> recently indicated that he was unable to observe a latent heat associated with the smectic-A–nematic transition and that the transition was presumably second order. Figure 1 shows a differential thermal analysis of the transition made on 9.1 mg of purified CBOOA at constant pressure. An apparent




FIG. 1. A differential thermal analysis of purified CBOOA. The sign of the peak changes sign with  $\Delta T$  indicating that the transition is not pure second order. It may be a small first-order transition or a  $\lambda$  transition. The absolute temperature calibration is  $\pm 1^\circ\text{C}$ . The temperature scale is in degrees Celsius.

latent heat is observed; however, the curve is not symmetric as a usual latent heat curve is, e.g., the solid-smectic transition shown to the right. Furthermore, there is a suggestion that one can amplify this bump by scanning the temperature faster. The peak begins to look like a specific heat jump when scanned rapidly; however, upon cooling, the bump becomes a dip which is evidence that the transition is not of pure second order. As purity deteriorated, the peak became smaller and smaller. In the impure limit it is virtually impossible to detect. The area shown in Fig. 1 would correspond to a latent heat of about 0.06 cal/g. The variation of this number from one curve to another is within the overall experimental error ( $\pm 20\%$ ). The lack of symmetry of the transition, the switching of the effect with  $\Delta T$ , and the apparent dependence of the effect on scanning rate suggest that it may be fruitful to investigate the possibility that the transition is a  $\lambda$  transition using more refined techniques. In very clean samples, however,  $T_c > 83.4^\circ\text{C}$ , we have observed a hysteresis effect in  $T_c$ ,  $\sim 0.2\text{--}0.4^\circ\text{C}$ . This would seem to indicate that the transition is first order.

The CBOOA used in the bend measurements was always recrystallized. The method<sup>5</sup> used was to dissolve 10% CBOOA in heptane, filter out a first small crop, reboil the liquid, filter out a second small crop, reboil, then finally reap the third crop  $\sim 50\%$  of the original amount. The original crystals were dark yellow. The yellow color fades in successive crops so that in the absolutely pure limit, the crystals should be white.<sup>5</sup> So far, the crystals have never been pure white but were considerably less yellow than the original crystals.

The methods used to measure the elastic constant of bend have been discussed elsewhere.<sup>6</sup> The technique consists of measuring the critical field of the well-known Fredericksz<sup>7</sup> transition by conoscopic observation of distortions induced by a magnetic field on monocrystalline samples at constant temperature. Thus, actually, one measures a threshold field whose square is proportional to  $K_3/\chi_a$ ,  $\chi_a$  being the anisotropy in the diamagnetic susceptibility. While measuring  $K_3/\chi_a$  one also finds<sup>8</sup>  $K_1/\chi_a$  (the splay constant which does not diverge) and the birefringence. In the following, the results for  $K_1/\chi_a$  are not presented, and I have not taken into account the temperature dependence of  $\chi_a(T)$ . A measure of the temperature dependence of  $\chi_a$  is given by the change in birefringence with temperature. This was found to be rather small ( $\sim 0.002/^\circ\text{C}$ ). A nonmagnetic Mettler hot stage was used to control the temperature.

Good monocrystalline samples are required for these measurements. The homeotropic alignment required for the  $K_3$  (bend) measurement was fabricated by coating the microscope slide and cover slip with a suitable surfactant.<sup>9</sup> Some of the samples were made by sealing the glass plates together with double bonding Mylar. This ensured that the sample thickness remained constant—even upon crystallization. Others were made by separating the glass plates with ordinary Mylar and the cell was sealed completely with rapid bonding epoxy. The Mylar samples showed no change in  $T_c$  even though the samples were kept several weeks. The measured values of  $K_3(T)$  were also highly reproducible from day to day and even from month to month.

The results for the bend measurements are shown in Fig. 2(a). Two runs are shown.  $T_c^*$  was determined by finding the best fit for several fixed values of  $T_c$ .  $\chi^2$  versus  $T_c$  is shown in Fig. 2(b). The  $T_c^*$  that best fits the data was chosen to be where the minimum occurred in the  $\chi^2$ . Thus the error shown for  $\gamma_3$  is a standard error based on a three-parameter fit [ $\gamma_3$ ,  $(K_3)_0$ , and  $(\text{const})_3$ ] but the fit is a four-parameter fit. The main result is that for the “clean” sample  $\gamma_3 = 0.52 \pm 0.03$ . This is a classical result. For the “dirty” sample,  $\gamma_3 = 0.60 \pm 0.01$ . The trend is that the lower the  $T_c$ , the more the exponent deviates from the classical value of 0.5.

The best fit gave  $T_c^* \sim T_{c, \text{meas}}$ . In fact,  $T_c^*$  differs from the measured  $T_c$  (when cooling the sample) by  $\lesssim 0.02^\circ\text{C}$ . When fitting only points within  $3^\circ\text{C}$  of  $T_c$ ,  $\gamma_3$  is not sensitive to the value




FIG. 2. (a) Plot of  $K_3(T) - (K_3)_0$  in arbitrary units versus  $T - T_c^*$ . The criterion for picking  $T_c^*$  is shown in (b). For clarity, the dirtier sample (right-hand scale) is shown one decade above the cleaner one (left-hand scale). For the dirty sample  $(K_3)_0 = 0.3067 \pm 0.406$  and for the cleaner sample  $(K_3)_0 = 0.196 \pm 0.942$  in arbitrary units. The lines drawn are the calculated best fits. (b)  $\chi^2$  versus  $T_c$  showing minimum for the definition of  $T_c^*$  for the curves shown in (a).  $T_c$  shown in degrees Celsius. The  $\chi^2$  shown is the  $\chi^2$  which resulted when  $T_c$  was fixed and  $\gamma_3$ ,  $(K_3)_0$ , and  $(\text{const})_i$  were found which best fit the data for that particular value of  $T_c$ .

of  $(K_3)_0$ . This can be seen from their respective computed standard errors [Fig. 2(a)]. As more points are added to the fit the standard error for  $(K_3)_0$  decreases. It is, in fact, possible to continue the fit for the cleaner samples to very close to the nematic-isotropic transition, i.e.,  $\sim 100^\circ\text{C}$ , without changing  $\gamma_3$ <sup>10</sup> or the minimum in  $\chi^2$  versus  $T_c^*$  (Fig. 2).

The crucial point in evaluating errors for this kind of an experiment is the uncertainty in  $T_c^*$ . The errors quoted above assume that  $T_c^*$  is defined infinitely precisely. Assuming an uncertainty in  $T_c^*$  of  $\pm 0.01^\circ\text{C}$ ,  $\gamma_3(\text{clean}) = 0.52 \pm 0.05$ . For an uncertainty of  $\pm 0.02^\circ\text{C}$ ,  $\gamma_3(\text{clean}) = 0.52 \pm 0.1$ . The  $\chi^2$  at  $T_c^* \pm 0.02^\circ\text{C}$  is somewhat more than twice its value at  $T_c^*$ . This tells us that  $T_c^*$ , determined in this way, can be relied upon to better than  $\pm 0.02^\circ\text{C}$  but not as close as  $\pm 0.01^\circ\text{C}$ . In order to define  $T_c^*$  better we require points closer to  $T_c^*$ . We are unable to continue these measurements indefinitely because very close to  $T_c^*$  the low-numbered birefringence fringes are not well defined.

Critical exponents are hypothesized to be universal functions of two parameters,<sup>11</sup> the dimensionality,  $d$ , of the system and the number,  $n$ , of components of the order parameter. Recently, Wilson<sup>12</sup> has calculated in the vicinity of small  $\delta = 4 - d$  an expansion for the exponents  $\gamma$  and  $\eta$  in terms of  $n$  and  $d$ . The exponent for the correlation length,  $\xi$ , is deduced from these two using

the Fisher equality,  $\nu = \gamma / (2 - \eta)$ , i.e.,

$$\xi = \xi_0 (T_c / |T - T_c|)^\nu, \quad (2)$$

where  $\xi_0$  is the zero-temperature correlation length. It is expected<sup>1</sup> that  $\delta K_3 \sim \xi_{\parallel}$ , where  $\xi_{\parallel}$  is the coherence length along the molecule as measured by McMillan using x rays.<sup>4</sup>  $\xi_{\perp}$  is the coherence length perpendicular to the molecule. Thus  $\gamma_3 \equiv \nu_{\parallel}$ , where  $\nu_{\parallel}$  is the exponent for  $\xi_{\parallel}$ .

Taking  $n = 2$  (for a complex order parameter) and  $d = 3$ , de Gennes estimates from the Wilson expansion that in the temperature region where the Landau theory fails,  $\nu = 0.66$ . However, McMillan finds<sup>4</sup>  $\xi_{\parallel} \sim (T - T_c)^{-0.75 \pm 0.01}$  but  $\xi_{\perp} \sim (T - T_c)^{-0.625 \pm 0.01}$  which indicates that either the simple Wilson model is inapplicable to the present system or the experimental errors are larger than have been stated. In fact, a recent calculation by Halperin, Lubensky, and Ma<sup>13</sup> indicates that there should always be a first-order nematic-smectic-A transition. The various exponents, therefore, would only have an approximate meaning and would depend upon the temperature range over which the fit is made.

De Gennes's calculation<sup>1</sup> is mean field. An estimate of the temperature range,  $\epsilon = (T - T_c) / T_c$ , where mean field theory fails,  $\epsilon \ll \epsilon_c$ , is given by the Ginzburg criterion,<sup>14</sup>

$$\epsilon_c = \frac{1}{32\pi^2} \left( \frac{k}{\Delta C_p} \frac{1}{(\xi_0)_{\perp}^2 (\xi_0)_{\parallel}} \right)^2, \quad (3)$$

where  $k$  is Boltzmann's constant and  $\Delta C_p$  is the specific heat jump at the transition.  $(\xi_0)_\perp = 3 \text{ \AA}$ ,  $(\xi_0)_\parallel = 10 \text{ \AA}$ ,<sup>4</sup> and from an extrapolation of Fig. 1,  $\Delta C_p \sim 0.09 \text{ cal/g } ^\circ\text{C}$ . Interpreting Fig. 1 as a specific heat jump gives  $\Delta C_p \sim 0.2\text{--}0.3 \text{ cal/g } ^\circ\text{C}$ . Using the smaller  $\Delta C_p \sim 0.09 \text{ cal/g } ^\circ\text{C}$ ,  $\epsilon_c \sim 5 \times 10^{-4}$ . Thus, only for  $(T - T_c) \ll 0.2^\circ\text{C}$  is non-classical behavior expected. This estimate may be broadened by impurities for then the  $\xi_0$ 's are smaller still.  $\Delta C_p$  is hard to define for a first-order transition. We think our estimate is an underestimate. It may also be broadened when one includes the fluctuations in the director field.

We recall that the x-ray measurements of McMillan, by definition, measure  $\xi_\parallel$  and  $\xi_\perp$  always, even in the nonclassical region. It has been shown that  $\delta K_3 \sim \xi_\parallel$  in the linearized, mean-field limit. It is to be noted that our dirtier sample does not exclude the possibility of 0.66, whereas our clean one does. We can broaden our nonclassical region by purposely adding controlled amounts of impurity to CBOOA. This results<sup>10</sup> in  $\gamma_3 \rightarrow 1.0$  (independent of impurity concentration) in agreement with previous results of Cheung and Meyers<sup>15</sup> on a compound known to be first order.

Although we cannot exclude completely the possibility that the nematic-smectic-A transition is a  $\lambda$  transition, we have anticipated the eventuality that it is a small first-order transition. Our data have been subjected to a least-squares fit. It appears possible to use the data to define  $T_c^*$  to  $\pm 0.015^\circ\text{C}$ . Our results are as follows: For the "clean" samples,  $\gamma_3$  approaches classical values. For the "dirty" sample,  $\gamma_3 = 0.6 \pm 0.08$ . As impurity is added, and hence  $T_c^*$  lowered,  $\gamma_3 \rightarrow 1$ . The Ginzburg criterion indicates that classical values are expected for clean samples except within a rather narrow temperature range near  $T_c$ .

A more complete account of this work, including the  $K_2(T)$  measurements and measurements on  $K_3(T)$  in the dirty limit, is in preparation.

I am particularly grateful to Patrick Lee and B. Halperin for many stimulating and fruitful discussions. I would also like to acknowledge stimulating discussions with H. E. Bair, D. Berreman, J. M. Daniels, R. Hewitt, A. Hutson, F. Kahn, and S. Meiboom.

<sup>1</sup>P. G. de Gennes, *Solid State Commun.* **10**, 753 (1972).

<sup>2</sup>A. Schmid, *Phys. Rev.* **180**, 527 (1969).

<sup>3</sup>F. C. Frank, *Discus. Faraday Soc.* **25**, 19 (1958).

<sup>4</sup>W. L. McMillan, *Phys. Rev. A* **7**, 1419 (1973).

McMillan refers to this compound as *n-p*-cyanobenzilidene *p*-amino-octyloxybenzene (CBAOB). According to G. N. Taylor (private communication), CBOOA corresponds to a terminology adopted by a more recent chemical convention so it is used here. CBAOB  $\equiv$  CBOOA. It is possible that McMillan's sample was slightly impure so that the values for  $(\xi_0)_\perp$  and  $(\xi_0)_\parallel$  are somewhat on the small side.

<sup>5</sup>Taylor, Ref. 4. It appears that the art of recrystallization depends upon guessing at what the impurities might be (in this case an "azo-compound," whence the yellowness of the crystals), then choosing a solvent which one thinks dissolves the impurity better than the compound (CBOOA) so that the crystals that form will be less rich in this impurity. I thank G. N. Taylor for sharing with me his expertise on the homologous series of this compound and for his assistance on the differential thermal analysis of purified CBOOA.

<sup>6</sup>C. E. Williams and P. E. Cladis, *Solid State Commun.* **10**, 357 (1972).

<sup>7</sup>V. Freedericksz and V. Zolina, *Trans. Faraday Soc.* **29**, 919 (1933).

<sup>8</sup>A. Saupe, *Z. Naturforsch.* **15a**, 815 (1960). For a more recent review, see H. Gruler, T. J. Scheffer, and G. Meier, *Z. Naturforsch.* **27a**, 966 (1972).

<sup>9</sup>F. J. Kahn, G. N. Taylor, and H. Schonhorn, to be published; F. J. Kahn, *Appl. Phys. Lett.* **22**, 386 (1973).

<sup>10</sup>P. E. Cladis, to be published.

<sup>11</sup>M. E. Fisher, *Rep. Progr. Phys.* **30**, 615 (1967).

<sup>12</sup>K. G. Wilson, *Phys. Rev. Lett.* **28**, 548 (1972).

<sup>13</sup>B. I. Halperin, T. Lubensky, and H. Ma, to be published.

<sup>14</sup>V. L. Ginzburg, *Fiz. Tverd. Tela* **2**, 2031 (1960)

[*Sov. Phys. Solid State* **2**, 1824 (1960)].

<sup>15</sup>L. Cheung and R. B. Meyers, *Phys. Lett.* **43A**, 261 (1973).